

APPENDICITIS CLINICAL PATHWAY

EXECUTIVE SUMMARY

Physician Owner(s): Dr. Zebulon Timmons



Primary Objective

Approximately one-third of all appendectomies at Children's Hospital & Medical Center are perforated (NSQIP definition). In 2018, we saw an increase in surgical site infections (SSI), *Streptococcus anginosus*, and readmissions for perforated appendicitis. In 2021-2022, Children's experienced increased variability in use of imaging in children for whom appendicitis was a concern, as well as an increased length of hospital stay.

The purpose of the Appendicitis Pathway is to standardize care and to decrease complications. The current departments that manage care for appendectomy and perforated appendectomy patients: Emergency Department, General Surgery and Operating Room, Inpatient, Outpatient Surgery Clinic, and Radiology. The intention of this pathway is to standardize patient evaluation for perforated appendicitis, expedite appropriate antibiotic treatments, reduce SSI rates, reduce return visits, and reduce return to the OR.

Emergency Department Recommendations

Inclusion Criteria

- Children 4 to 21 with clinical suspicion of appendicitis

Exclusion Criteria

*Acute appendicitis can occur in **any** of the following excluded populations, but likely require different considerations in evaluation.*

- Children <4 years
- Toxic appearance
- Hemodynamic instability
- Previous appendectomy
- Pregnancy
- History of IBD
- Trauma patients
- CT done at outside hospital
- Ongoing treatment for malignancy
- History of organ transplant
- Severe developmental delay

SCORING TOOLS

- Only 50-70% of patients with appendicitis are correctly diagnosed at initial assessment because there is no definitive sign, symptom, or laboratory value. The implementation and evolution of scoring systems have increased diagnostic accuracy from 58% to 71%.³¹
- The Pediatric Appendicitis Score (PAS) is a scoring system for children 3 years to 18 years and generates a risk score based on eight statistically significant variables. Variables include findings from the patient's history, physical, and laboratory results.³¹
- PAS serves as a clinical decision support tool to guide a clinician in determining whether to observe or operate.³¹

APPENDICITIS CLINICAL PATHWAY

EXECUTIVE SUMMARY

Physician Owner(s): Dr. Zebulon Timmons

- PAS has been validated in multicenter studies; one study showed a sensitivity of 1, specificity of 0.92, positive predictive value of 0.96, and negative predictive value of 0.99.³¹

LABORATORY EVALUATION

- WBC <11,000 unlikely to be appendicitis in non-visualized and non-diagnostic ultrasounds with NPV 95% and 96% respectively.¹⁰
- A left shift with normal WBC is seen in early appendicitis.³⁹

DIAGNOSTIC IMAGING

- Ultrasound (US) has emerged as the preferential first-line modality for imaging in suspected appendicitis.¹²
- Despite its advantages,¹² US by itself has limitations in diagnosing appendicitis²⁸, with the rate of equivocal examinations in some studies approaching 50%.^{29, 30, 34, 36}
Category 0-4 definitions for ultrasound reads to guide diagnosis and initiation of antibiotic therapy:
 - Category 0:** Examination nondiagnostic/poor visualization due to habitus. Further imaging is to be based on full clinical evaluation.
 - Category 1:** Appendix seen and normal.
 - Category 2:** Appendix not seen or not fully seen, but no specific secondary signs of appendicitis.
 - Category 3:** Appendix not seen or not fully seen but concerning RLQ findings/inflammation. Additional imaging may be beneficial.
 - Category 4:** Positive exam for appendicitis.
- Several scoring systems, including the pediatric appendicitis score (PAS),³¹ can help the clinician with diagnosis, but also do not perform well in isolation at diagnosing appendicitis.
4-5, 11, 14-15, 23-24, 32
- An intravenous contrast-enhanced computed tomography (CT) scan has historically been utilized in children with equivocal screening US results.¹⁸
- Due to rising awareness of a potential CT-related risk of radiation-induced malignancies in children,^(7, 27, 38) there has been a decrease in the CT rate over the past decade.³
- Magnetic Resonance Imaging (MRI) has emerged as an accurate second-line modality to US with good diagnostic accuracy and quick protocols that do not require sedation.¹³
- Children's will trial utilizing the same scoring system for MRI Appendicitis exams as it currently uses for US studies (see above) to help guide decision-making.
- There has also been evidence that serial exams and repeat US can help avoid many CT scans and their associated radiation.³³
- In summary, when appendicitis is suspected an US should be obtained and when equivocal an MRI and/or serial exam should be employed to avoid CT whenever possible.**

ANTIMICROBIAL THERAPY

- Otherwise, healthy children without cephalosporin allergy AND without suspicion for perforation, abscess or phlegmon should be started on cefoxitin. Cefoxitin covers MSSA, *Streptococcus anginosus*, enteric gram negatives and some anaerobic bacteria. It does not cover *Pseudomonas aeruginosa*; however, regimens without pseudomonal coverage have been shown equally efficacious in pediatric appendicitis as regimens with anti-pseudomonal drugs^{8,19, 26} including patients with complicated appendicitis.^{17, 21-22}
- Otherwise, healthy children without suspicion of perforation, abscess or phlegmon but who have cephalosporin allergies should be started on levofloxacin plus metronidazole.³⁵
- In patients with suspicion for or confirmation of perforation, abscess or phlegmon, therapy should be initiated with piperacillin/tazobactam. In recent years, perforated appendicitis has been increasingly complicated by late abscesses due to *Streptococcus anginosus*, which piperacillin/tazobactam covers. For patients with penicillin allergy, consider levofloxacin plus metronidazole or ceftriaxone plus metronidazole.
- Immunocompromised patients or patients with concern for sepsis or severe disease should be started on piperacillin/tazobactam unless CNS disease cannot be excluded, in which case they should be started on cefepime + metronidazole. Infectious Disease consultation is recommended.
- The optimal time for administration of preoperative doses is within 60 minutes before surgical incision to reduce the rate of surgical site infection (SSIs) based on the Clinical Practice Guidelines for Antimicrobial Prophylaxis in Surgery.⁶
- Surgical Site infections were significant lower when preoperative antibiotics were given between 16-60 minutes before incision compared to 0-15 minutes and 121-180 minutes prior to incision. SSI risk was higher for patients receiving infusions 61-120 minutes before incision (confidence interval 0.98-5.61).⁶
- Ideally, an antimicrobial agent for surgical prophylaxis should (1) prevent SSI, (2) prevent SSI-related morbidity and mortality, (3) reduce the duration and cost of health care (when the costs associated with the management of SSI are considered, the cost-effectiveness of prophylaxis becomes evident)^{1,2}, (4) produce no adverse effects, and (5) have no adverse consequences for the microbial flora of the patient or the hospital.³⁷ To achieve these goals, an antimicrobial agent should be (1) active against the pathogens most likely to contaminate the surgical site, (2) given in an appropriate dosage and at a time that ensures adequate serum and tissue concentrations during the period of potential contamination, (3) safe, and (4) administered for the shortest effective period to minimize adverse effects, the development of resistance, and costs.^{16, 20, 25}

Metrics

Emergency Department

1. Increase proportion of patients who receive IV Antibiotics prior to Pre-op dose (antibiotics given 0-60mins prior to incision count as bundle dose; doses received 60+ mins prior to incision count as prior to Pre-op dose) by 10% from baseline by September 2023. – Outcome Metric

APPENDICITIS CLINICAL PATHWAY

EXECUTIVE SUMMARY

Physician Owner(s): Dr. Zebulon Timmons



- a. Denominator = all appendicitis patients with appendectomy procedure listed on OR schedule
2. Decrease proportion of total appendicitis patients who undergo an Abdominal CT during evaluation in the CHMC ED by 10% from baseline by September 2023. – Outcome Metric
 - a. Denominator = any patient with RLQ US or MR appendicitis WO contrast ordered in CHMC ED
3. Increase utilization of ED Appendicitis order set for evaluation of patients diagnosed with appendicitis to 50% by September 2023. – Process Metric
 - a. Denominator = any patient with RLQ US or MR appendicitis WO contrast ordered in CHMC ED
4. Monitor ED LOS (minutes) from triage to disposition. – Balancing Metric
 - a. Denominator = any patient with RLQ US or MR appendicitis WO contrast ordered in CHMC ED

Rationale

Appendicitis is the most common surgical emergency in pediatrics. Despite the frequency with which it is seen, appendicitis can still be elusive, as it presents in a myriad of different ways. The diagnostic uncertainty which comes with appendicitis has led to variability in care across different institutions and often between providers at a single site.

The purpose of a clinical pathway for appendicitis is to reduce these inconsistent approaches. It will allow for focused care based on the extensive available evidence on how to appropriately diagnose and manage this condition. Following the best available evidence will lead to:

1. A reduction in unnecessary imaging, especially modalities that expose patients to ionizing radiation.
2. Better timing of antibiotic administration as well as more focused antibiotic coverage with a focus on stewardship.
3. Reduced confusion for families, staff, and providers as we are all involved are following the same plan.

A clinical pathway like this is often borne out of compromise, so some who have practiced differently in the past may be guided towards different practices that will require education and monitoring for compliance. We will also have to monitor patient outcomes to remain confident that these interventions continue to be the best choice for our specific patient population.

Team Members

Champion: Zebulon Timmons, M.D. (Emergency Department Division Chief)

Members:

- Jennifer Wang, D.O. (Emergency Department Medical Director)
- Elizabeth Stenvers PA-C (Emergency Department)
- Abdalla Zarroug, M.D. (General Surgery Division Chief)
- Angela Hanna, M.D. (General Surgery)
- Patrick Thomas, M.D. (General Surgery)
- Robert Cusick, M.D. (General Surgery)

APPENDICITIS CLINICAL PATHWAY

EXECUTIVE SUMMARY

Physician Owner(s): Dr. Zebulon Timmons

- Megan Fuller, M.D. (General Surgery)
- Lauren Maskin, M.D. (Medical Director for Inpatient Medical-Surgical Services)
- Alice Sato, M.D., PhD (Infectious Disease)
- Susie Beedle, RN (Infection Prevention)
- Sandra Allbery, M.D. (Radiology)
- Jennifer Zwiener, PharmD (Pharmacy & Antimicrobial Stewardship)
- Kelsey Spackler DNP APRN-NP CPNP-AC/PC (Clinical Effectiveness)
- Abby Vipond, MSN, APRN, FNP-C, (Clinical Effectiveness)
- Ellen Kerns, PhD, MPH (Care Transformation Business Intelligence)

Evidence

1. Alerany C, Campany D, Monerde J, et al. Impact of local guidelines and an integrated dispensing system on antibiotic prophylaxis quality in a surgical centre. *J Hosp Infect.* 2005; 60:111–7.
2. Allerberger F, Gareis R, Jindrák V, et al. Antibiotic stewardship implementation in the EU: the way forward. *Expert Rev Anti Infect Ther.* 2009; 7:1175–83.
3. Bachur RG, Hennelly K, Callahan MJ, Monuteaux MC. Advanced radiologic imaging for pediatric appendicitis, 2005–2009: trends and outcomes. *J Pediatrics* 2012;160:1034–8.
4. Bhatt M, Joseph L, Ducharme FM, Dougherty G, McGillivray D. Prospective validation of the pediatric appendicitis score in a Canadian pediatric emergency department. *Acad Emerg Med* 2009; 16:591–6.
5. Bond GR, Tully SB, Chan LS, Bradley RL. Use of the MANTRELS score in childhood appendicitis: a prospective study of 187 children with abdominal pain. *Ann Emerg Med* 1990;19:1014–8.
6. Bratzler DW, Dellinger EP, Olsen KM, Perl TM, Auwaerter PG, Bolon MK, Fish DN, Napolitano LM, Sawyer RG, Slain D, Steinberg JP, Weinstein RA; American Society of Health-System Pharmacists; Infectious Disease Society of America; Surgical Infection Society; Society for Healthcare Epidemiology of America. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Am J Health Syst Pharm.* 2013 Feb 1;70(3):195-283. doi: 10.2146/ajhp120568. PMID: 23327981.
7. Brenner D, Elliston C, Hall E, Berdon W. Estimated risks of radiation-induced fatal cancer from pediatric CT. *Am J Roentgenol* 2001;176:289–96.
8. Cameron, Danielle B. MD, MPH*; Melvin, Patrice MPH†; Graham, Dionne A. PhD†; Glass, Charity C. MD, MPP*; Serres, Stephanie K. MD, PhD*; Kronman, Matthew P. MD, MSCE‡; Saito, Jacqueline M. MD, MSCI§; Rangel, Shawn J. MD, MSCE* Extended Versus Narrow-spectrum Antibiotics in the Management of Uncomplicated Appendicitis in Children, *Annals of Surgery*: July 2018 - Volume 268 - Issue 1 - p 186-192 doi: 10.1097/SLA.0000000000002349
9. Child J, Chen X et al. Pharmacokinetic and Pharmacodynamic properties of metronidazole in pediatric patients with acute appendicitis: A prospective study. *JPIDS* 2017; 2018;00(00):1–6. DOI: 10.1093/jpids/piy040
10. Cohen B, Bowling J, Midulla P, Shlasko E, Lester N, Rosenberg H, Lipskar A. The non-diagnostic ultrasound in appendicitis: is a non-visualized appendix the same as a negative

APPENDICITIS CLINICAL PATHWAY

EXECUTIVE SUMMARY

Physician Owner(s): Dr. Zebulon Timmons



- study? *J Pediatr Surg.* 2015 Jun;50(6):923-7. doi: 10.1016/j.jpedsurg.2015.03.012. Epub 2015 Mar 17. PMID: 25841283.
11. Dingemann J, Ure B. Imaging and the use of scores for the diagnosis of appendicitis in children. *Eur J Pediatr Surg* 2012;22:195–200.
 12. Doria AS, Moineddin R, Kellenberger CJ, et al. US or CT for diagnosis of appendicitis in children and adults? A meta-analysis. *Radiology* 2006;241:83–94.
 13. Eng KA, Abadeh A, Ligocki C, Lee YK, Moineddin R, Adams-Webber T, Schuh S, Doria AS. *Radiology.* 2018 Sep;288(3):717-727. doi: 10.1148/radiol.2018180318. Epub 2018 Jun 19. PMID: 29916776
 14. Escriba A, Gamell AM, Fernandez Y, Quintilla JM, Cubells CL. Prospective validation of two systems of classification for the diagnosis of acute appendicitis. *Pediatr Emerg Care* 2011;27:165–9.
 15. Goldman RD, Carter S, Stephens D, Antoon R, Mounstephen W, Langer JC. Prospective validation of the pediatric appendicitis score. *J Pediatrics* 2008;153:278–82.
 16. Gorbach SL, Condon RE, Conte JE Jr, et al. Evaluation of new anti-infective drugs for surgical prophylaxis. *Clin Infect Dis.* 1992; 15(suppl 1):S313–38.
 17. Hamdy RF, Handy LK, Spyridakis E, Dona D, Bryan M, Collins JL, Gerber JS. Comparative Effectiveness of Ceftriaxone plus Metronidazole versus Anti-Pseudomonal Antibiotics for Perforated Appendicitis in Children. *Surg Infect (Larchmt).* 2019 Jul;20(5):399-405. doi: 10.1089/sur.2018.234. Epub 2019 Mar 15. PMID: 30874482; PMCID: PMC6555178.
 18. Howell JM, Eddy OL, Lukens TW, Thiessen ME, Weingart SD, Decker WW. Clinical policy: critical issues in the evaluation and management of emergency department patients with suspected appendicitis. *Ann Emerg Med* 2010;55:71–116.
 19. Hurst AL, Olson D, Somme S, Child J, Pyle L, Ranade D, Stamatoiu A, Crombleholme T, Parker SK. Once-Daily Ceftriaxone Plus Metronidazole Versus Ertapenem and/or Cefoxitin for Pediatric Appendicitis. *J Pediatric Infect Dis Soc.* 2017 Mar 1;6(1):57-64. doi: 10.1093/jpids/piv082. PMID: 26703242.
 20. Källman J, Friberg Ö. Antibiotic prophylaxis in cardiac surgery—general principles. *APMIS.* 2007;115:1012–5.
 21. Kashtan MA, Graham DA, Melvin P, Hills-Dunlap JL, Anandalwar SP, Rangel SJ. Ceftriaxone with metronidazole versus piperacillin/tazobactam in the management of complicated appendicitis in children: Results from a multicenter pediatric NSQIP analysis. *J Pediatr Surg.* 2021 Nov 20:S0022-3468(21)00784-3. doi: 10.1016/j.jpedsurg.2021.11.009. Epub ahead of print. PMID: 34876294.
 22. Kronman MP, Oron AP, Ross RK, Hersh AL, Newland JG, Goldin A, Rangel SJ, Weissman SJ, Zerr DM, Gerber JS. Extended- Versus Narrower-Spectrum Antibiotics for Appendicitis. *Pediatrics.* 2016 Jul;138(1):e20154547. doi: 10.1542/peds.2015-4547. PMID: 27354453.
 23. Kulik DM, Uleryk EM, Maguire JL. Does this child have appendicitis? A systematic review of clinical prediction rules for children with acute abdominal pain. *J Clin Epidemiol* 2013;66:95–104.
 24. Mandeville K, Pottker T, Bulloch B, Liu J. Using appendicitis scores in the pediatric ED. *Am J Emerg Med* 2011;29:972–7.
 25. Mangram AJ, Horan TC, Pearson ML, et al. Guideline for prevention of surgical site infection. *Infect Control Hosp Epidemiol.* 1999; 20:250–78.

APPENDICITIS CLINICAL PATHWAY

EXECUTIVE SUMMARY

Physician Owner(s): Dr. Zebulon Timmons



26. Paek H, Pham H, Gagliardo C, Caruso-Prendergast P. Cefoxitin vs. Piperacillin/Tazobactam for Acute Appendicitis in Pediatric Patients. *Open Forum Infect Dis*. 2017;4(Suppl 1):S500. Published 2017 Oct 4. doi:10.1093/ofid/ofx163.1295
27. Pearce MS, Salotti JA, Little MP, et al. Radiation exposure from CT scans in childhood and subsequent risk of leukaemia and brain tumours: a retrospective cohort study. *Lancet* 2012;380:499–505.
28. Pena BM, Taylor GA. Radiologists' confidence in interpretation of sonography and CT in suspected pediatric appendicitis. *Am J Roentgenol* 2000;175:71–4.
29. Ramarajan N, Krishnamoorthi R, Barth R, et al. An interdisciplinary initiative to reduce radiation exposure: evaluation of appendicitis in a pediatric emergency department with clinical assessment supported by a staged ultrasound and computed tomography pathway. *Acad Emerg Med* 2009;16:1258–65.
30. Saito JM, Yan Y, Evashwick TW, Warner BW, Tarr PI. Use and accuracy of diagnostic imaging by hospital type in pediatric appendicitis. *Pediatrics* 2013;131:e37–44.
31. Samuel M. Pediatric appendicitis score. *J Pediatr Surg* 2002;37:877–81.
32. Schneider C, Kharbanda A, Bachur R. Evaluating appendicitis scoring systems using a prospective pediatric cohort. *Ann Emerg Med* 2007;49:778–84.
33. Schuh S, Chan K, Langer JC, Kulik D, Preto-Zamperlini M, Aswad NA, Man C, Mohanta A, Stephens D, Doria AS. Properties of serial ultrasound clinical diagnostic pathway in suspected appendicitis and related computed tomography use. *Acad Emerg Med*. 2015 Apr;22(4):406-14. doi: 10.1111/acem.12631. Epub 2015 Mar 24. PMID: 25808065.
34. Schuh S, Man C, Cheng A, et al. Predictors of nondiagnostic ultrasound scanning in children with suspected appendicitis. *J Pediatr* 2011;158:112–8.
35. St Peter SD, Tsao K, Spilde TL, et al. Single daily dosing cefTRIAXone and metronidazole vs standard triple antibiotic regimen for perforated appendicitis in children: a prospective randomized trial. *J Pediatr Surg* 2008;43:981-5.
36. Trout AT, Sanchez R, Ladino-Torres MF, Pai DR, Strouse PJ. A critical evaluation of US for the diagnosis of pediatric acute appendicitis in a real-life setting: how can we improve the diagnostic value of sonography? *Pediatr Radiol* 2012;42:813–23.
37. Voit SB, Todd JK, Nelson B, et al. Electronic surveillance system for monitoring surgical antimicrobial prophylaxis. *Pediatrics*. 2005; 116:1317–22.
38. Wan MJ, Krahn M, Ungar WJ, et al. Acute appendicitis in young children: cost-effectiveness of US versus CT in diagnosis—a Markov decision analytic model. *Radiology* 2009;250:378–86.
39. Wang LT, Prentiss KA, Simon JZ, Doody DP, Ryan DP. The use of white blood cell count and left shift in the diagnosis of appendicitis in children. *Pediatr Emerg Care*. 2007 Feb;23(2):69-76. doi: 10.1097/PEC.0b013e31802d1716. PMID: 17351404.