

APPENDICITIS PATHWAY

EMERGENCY DEPARTMENT


Exclusion Criteria:

- Children < 4 years
- Toxic appearance
- Hemodynamic instability
- Neurologic changes
- Previous appendectomy
- Pregnancy
- History of IBD
- Trauma patients
- CT done at OSH
- Ongoing treatment for malignancy
- History of organ transplant
- Severe developmental delay

Acute appendicitis can occur in ANY of above excluded patient populations, but likely require different considerations in evaluation


Imaging Grading:

- Category 0:** Examination nondiagnostic/poor visualization due to habitus
- Category 1:** Appendix seen and normal
- Category 2:** Appendix not fully seen but no secondary signs of appendicitis
- Category 3:** Appendix not seen or not fully seen but concerning RLQ findings/ inflammation
- Category 4:** Positive for appendicitis



APPENDICITIS PATHWAY

EMERGENCY DEPARTMENT



Patient Risk Assessment [‡]	
High probability acute appendicitis	Symptoms <48hrs
	Migration of pain from periumbilical region to RLQ
	Anorexia, nausea, vomiting
	Pain preceding vomiting
	Pain w ith movement (cough, car ride, jumping, heel tap)
Equivocal acute appendicitis	RLQ tenderness w ith or w ithout tenderness
	Presenting w ith focal abdominal tenderness (usually right sided) w ith some of the features of high probability acute appendicitis
Suspicious for complex appendicitis (perforation/abscess)	Systemic toxicity (also exclusion criteria)
	Fever
	Prolonged symptoms >48-72hrs
	Urinary or rectal urgency
	Palpable RLQ mass
	WBC, ANC, CRP consistent w ith marked inflammation

Pediatric Appendicitis Score [◊]	
Low Risk <4; High Risk ≥7	
Nausea/vomiting	1
Anorexia	1
Migration of pain to RLQ	1
Fever	1
Pain w ith movement	2
RLQ tenderness	2
WBC >10,000	1
ANC >7,500	1
<i>An experienced physician's clinical impression performs comparably to this score</i>	

APPENDICITIS PATHWAY

EMERGENCY DEPARTMENT

Preoperative Antibiotics[®]

Indicated for ALL patients with Acute Appendicitis (to be given at time of category 4 diagnosis or in discussion with Surgeon)

Healthy non-allergic patients without suspicion of perforation, abscess or phlegmon	Cefoxitin IV 40mg/kg/dose (max 2,000mg) (Must be adjusted in renal impairment)
Healthy patients without suspicion of perforation, abscess or phlegmon with cephalosporin allergy	Levofloxacin IV 6 month to < 5 years: 10mg/kg/dose BID (max 750mg) ≥ 5 years: 10mg/kg once daily (max 750mg) AND Metronidazole IV 30mg/kg/dose once daily (max 1,500mg)
Patients with suspicion of perforation, abscess or phlegmon	Piperacillin/Tazobactam IV 100mg/kg/dose (max 3,000mg) (Must be adjusted in renal impairment)
Patients with suspicion of perforation, abscess or phlegmon with severe penicillin allergy	Levofloxacin IV 6 month to < 5 years: 10mg/kg/dose BID (max 750mg) ≥ 5 years: 10mg/kg once daily (max 750mg) AND Metronidazole IV 30mg/kg/dose once daily (max 1,500mg)
Patients who are immunocompromised or very ill	Piperacillin/Tazobactam IV 100mg/kg/dose (max 3,000mg) (Consider Infectious Disease consult)

Laboratory Evaluation[^]

All patients May choose to defer blood tests in patients without other indications for PIV	CBC with ANC ¹
	CRP ²
	UA ³
As clinically indicated	CMP
	Lipase, amylase
	Urine HCG (all female patients >11yrs/or menstruating females)
	Urine GC/Chlamydia, vaginal sw ab for Trichomonas
Ill patients	Blood Culture
	Coagulation Studies
	Lactate
	Procalcitonin
	Type and Screen (not indicated for routine appendectomy)

^{1,2}WBC >11,000 **AND/OR** CRP >1.0 is found in 95% of non-perforated appendicitis and 100% of perforated, abscess or phlegmon appendicitis

Therefore, if both are normal, consider an alternate diagnosis

If one or the other is elevated, it adds little value to the diagnostic evaluation beyond PAS score

²UA in appendicitis may occasionally demonstrate sterile pyuria

Leukouria without bacteria or nitrates present should not dissuade one from the diagnosis of appendicitis